

 spncrfx@gmail.com

 @foxandtheflu

Introduction to Probability and Likelihood in R

Spencer Fox
14 October 2016

Week 6 goals

Week 6 goals

- Working with probability distributions

Week 6 goals

- Working with probability distributions
- Likelihood
 - Calculating likelihoods
 - Interpreting likelihood results

R probability distribution functions
are extremely consistent

R probability distribution functions
are extremely consistent

xdist()


R probability distribution functions
are extremely consistent

xdist()

r, d, p, or q



R probability distribution functions
are extremely consistent



Drawing (r)andom (norm)al samples


```
rnorm(n = 10000, mean = 0, sd = 1)
```

Drawing (r)andom (norm)al samples


```
rnorm(n = 10000, mean = 0, sd = 1) %>%  
  hist()
```

Drawing (r)andom (norm)al samples

```
rnorm(n = 10000, mean = 0, sd = 1) %>%  
  hist()
```




Getting the (d)ensity or likelihood of a data point




Getting the (d)ensity or likelihood of a data point

```
dnorm(x = 2)
```


```
[1] 0.05399097
```




Getting cumulative (p)robability and the inverse (q)umulative probability




Getting cumulative (p)robability and the inverse (q)umulative probability




Getting cumulative (p)robability and the inverse (q)umulative probability




Getting cumulative (p)robability and the inverse (q)umulative probability




Getting cumulative (p)robability and the inverse (q)umulative probability



Getting cumulative (p)robability and the inverse (q)umulative probability



Getting cumulative (p)robability and the inverse (q)umulative probability



1st Programming Exercise

Likelihood is the probability of
your data under a given model

$$P(Y \mid \theta)$$


Likelihood is the probability of your data under a given model

$$P(Y \mid \theta)$$


Data



Likelihood is the probability of your data under a given model




We often denote the parameters
and not the model itself




Our first model: linear regression

$$Y = \beta X + a$$




Our first model: linear regression

$$Y = \beta X + a$$



Our first model: linear regression


$$Y_i \sim N(\beta X_i + a, \sigma)$$




Our first model: linear regression

$$Y_i \sim N(\beta X_i + a, \sigma)$$


assumed sd




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




Try a different model



$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$



`dnorm(x=Yi, mean= $\beta X_i + \alpha$, sd= σ)`


$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$



$$\text{dnorm}(x=Y_i, \text{mean}=\beta X_i + \alpha, \text{sd}=\sigma)$$

$$P(Y \mid \beta, \alpha) = \text{product for all } i$$

$$Y_i \sim N(\beta X_i + \alpha, \sigma)$$




`dnorm(x=Yi, mean= $\beta X_i + \alpha$, sd= σ , log=T)`

$P(Y \mid \beta, \alpha) = \text{sum}$ for all i


2nd Programming Exercise

```
function_name <- function(arg1, arg2, ...){  
  carry out function process  
  output  
}
```


Once we have likelihood function we
can systematically test parameters




Once we have likelihood function we can systematically test parameters




Once we have likelihood function we
can systematically test parameters




Once we have likelihood function we
can systematically test parameters




```
> lm(y~x, data =sim1)
```



```
> lm(y~x, data =sim1)
```





```
> lm(y~x, data =sim1)
```

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	4.2208	0.8688	4.858	4.09e-05	***
x	2.0515	0.1400	14.651	1.17e-14	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


```
> lm(y~x, data =sim1)
```

Call:

```
lm(formula = y ~ x, data = sim1)
```

Residuals:

	Min	1Q	Median	3Q	Max
	-4.1469	-1.5197	0.1331	1.4670	4.6516

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.203 on 28 degrees of freedom

Multiple R-squared: 0.8846, Adjusted R-squared: 0.8805

F-statistic: 214.7 on 1 and 28 DF, p-value: 1.173e-14

```
> lm(y~x, data =sim1)
```

should be ~ normal



Call:

```
lm(formula = y ~ x, data = sim1)
```

Residuals:

Min	1Q	Median	3Q	Max
-4.1469	-1.5197	0.1331	1.4670	4.6516

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.203 on 28 degrees of freedom

Multiple R-squared: 0.8846, Adjusted R-squared: 0.8805

F-statistic: 214.7 on 1 and 28 DF, p-value: 1.173e-14

```
> lm(y~x, data =sim1)
```

should be ~ normal

Call:

```
lm(formula = y ~ x, data = sim1)
```

MLE

Residuals:

Min	1Q	Median	3Q	Max
-4.1469	-1.5197	0.1331	1.4670	4.6516

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.203 on 28 degrees of freedom

Multiple R-squared: 0.8846, Adjusted R-squared: 0.8805

F-statistic: 214.7 on 1 and 28 DF, p-value: 1.173e-14

```
> lm(y~x, data =sim1)
```

should be ~ normal

Call:

```
lm(formula = y ~ x, data = sim1)
```

MLE

Standard error

Residuals:

Min	1Q	Median	3Q	Max
-4.1469	-1.5197	0.1331	1.4670	4.6516

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.203 on 28 degrees of freedom

Multiple R-squared: 0.8846, Adjusted R-squared: 0.8805

F-statistic: 214.7 on 1 and 28 DF, p-value: 1.173e-14

```
> lm(y~x, data =sim1)
```

should be ~ normal

Call:

```
lm(formula = y ~ x, data = sim1)
```

MLE

Standard error

Residuals:

Min	1Q	Median	3Q	Max
-4.1469	-1.5197	0.1331	1.4670	4.6516

coefficient t-statistic

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.203 on 28 degrees of freedom

Multiple R-squared: 0.8846, Adjusted R-squared: 0.8805

F-statistic: 214.7 on 1 and 28 DF, p-value: 1.173e-14


```
> lm(y~x, data =sim1)
```

should be ~ normal

```
Call:  
lm(formula = y ~ x, data = sim1)
```

MLE

Standard error

```
Residuals:
```

Min	1Q	Median	3Q	Max
-4.1469	-1.5197	0.1331	1.4670	4.6516

coefficient t-statistic

```
Coefficients:
```

p-val

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 2.203 on 28 degrees of freedom
```

```
Multiple R-squared:  0.8846,    Adjusted R-squared:  0.8805
```

```
F-statistic: 214.7 on 1 and 28 DF,  p-value: 1.173e-14
```

```
> lm(y~x, data =sim1)
```

should be ~ normal

Call:

```
lm(formula = y ~ x, data = sim1)
```

MLE

Standard error

Residuals:

Min	1Q	Median	3Q	Max
-4.1469	-1.5197	0.1331	1.4670	4.6516

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	4.2208	0.8688	4.858	4.09e-05 ***
x	2.0515	0.1400	14.651	1.17e-14 ***

coefficient t-statistic

p-val

fit

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.203 on 28 degrees of freedom

Multiple R-squared: 0.8846, Adjusted R-squared: 0.8805

F-statistic: 214.7 on 1 and 28 DF, p-value: 1.173e-14

3rd Programming Exercise